

A COMPARATIVE STUDY OF THE MASS DISTRIBUTION OF EXTREME ULTRAVIOLET-SELECTED WHITE DWARFS ¹

R. NAPIWOTZKI

Dr. Remeis-Sternwarte, Sternwartstr. 7, 96049 Bamberg, Germany
 email: *ai23@sternwarte.uni-erlangen.de*

PAUL J. GREEN

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138
 email: *pgreen@cfa.harvard.edu*

AND

REX A. SAFFER

Dept. of Astronomy & Astrophysics, Villanova University, 800 Lancaster Ave., Villanova, PA 19085
 email: *saffer@ast.vill.edu*

ApJ Vol. 517, May 20, 1999, in press

ABSTRACT

We present new determinations of effective temperature, surface gravity, and masses for a sample of 46 hot DA white dwarfs selected from the EUVE and ROSAT Wide Field Camera bright source lists in the course of a near-IR survey for low mass companions. Our analysis, based on hydrogen NLTE model atmospheres, provides a map of LTE correction vectors, which allow a thorough comparison with previous LTE studies. We find previous studies underestimate both the systematic errors and the observational scatter in the determination of white dwarf parameters obtained via fits to model atmospheres.

The structure of very hot or low mass white dwarfs depends sensitively on their history. To compute white dwarf masses, we thus use theoretical mass-radius relations that take into account the complete evolution from the main sequence. We find a peak mass of our white dwarf sample of $0.59M_{\odot}$, in agreement with the results of previous analyses. However, we do not confirm a trend of peak mass with temperature reported in two previous analyses.

Analogous to other EUV selected samples, we note a lack of low mass white dwarfs, and a large fraction of massive white dwarfs. Only one white dwarf is likely to have a helium core. While the lack of helium white dwarfs in our sample can be easily understood from their high cooling rate and therefore low detection probability in our temperature range, this is not enough to explain the large fraction of massive white dwarfs. This feature very likely results from a decreased relative sample volume for low mass white dwarfs caused by interstellar absorption in EUV selected samples.

Subject headings: stars: atmospheres — stars: evolution — stars: white dwarfs — binaries: close

1. INTRODUCTION

UV observations of EUV-detected stars have revealed the presence of about 15 hot white dwarf companions to bright stars in non-interacting binary systems (e.g., Burleigh, Barstow, & Fleming 1997). At optical wavelengths, these WDs are hidden because of their close proximity to much more luminous companions which are main sequence (spectral type K or earlier) or evolved stars.

A fascinating variety of objects are known or proposed to contain white dwarf (WD) stars in interacting binary systems. A partial list includes novae, cataclysmic variables, symbiotic stars, Ba and CH giants, Feige 24-type systems and dwarf carbon stars (Green & Margon 1994). These systems offer great insights to evolution and dynamical processes in binaries.

A number of interacting binary systems where the white dwarf is the primary (i.e., optically brightest) star have also been found among EUV-detected systems (e.g., 6 close, interacting white dwarf/red dwarf binaries by Vennes & Thorstensen 1994). Optical or ultraviolet spec-

tral observations are most commonly used to detect companions to WD primaries, by searching for (1) the presence of narrow Balmer line emission overlying the broad smooth Balmer absorption of the WD, (2) a composite white dwarf + main sequence spectrum, or (3) radial velocity (RV) variations. However, only WDs with very close, or intrinsically active companions will be found by method (1). For hot white dwarf systems, composite spectra (2) are only expected to be visible if the companion's spectral type is early enough. RV variations (3) require multiple observations at high spectral resolution, and detection strongly favors close and/or massive companions.

All the discoveries mentioned above have been strongly dominated by these selection effects, with companions biased to earlier types than predicted by the simulations of deKool & Ritter (1993) and others. Scaling from deKool & Ritter's results, Vennes & Thorstensen (1994) estimate that "at least twice as many close binary systems remain to be identified from EUV surveys, most of them with a low mass secondary." The resulting sample of binaries known

¹SPECTRAL OBSERVATIONS REPORTED HERE WERE OBTAINED WITH THE MULTIPLE MIRROR TELESCOPE, A JOINT FACILITY OF THE UNIVERSITY OF ARIZONA AND THE SMITHSONIAN INSTITUTION, AND WITH THE BOK TELESCOPE AT THE STEWARD OBSERVATORY OF THE UNIVERSITY OF ARIZONA.

to date therefore must diverge strongly from the intrinsic distribution, in overall normalization, as well as in mass and spectral type of the main sequence companions.

The current study, conceived as a complement to optical studies, began as a near-IR photometric survey for low mass companions to hot white dwarfs (WDs). By investigating only EUV-detected WDs, we obtain a very reasonably-sized but complete sample of young WDs, next to which very late-type dwarf companions can be detected in the near-infrared by searching for a K excess over that expected from the white dwarf. Many hot white dwarfs ($T_{\text{eff}} > 24000\text{K}$; Finley et al. 1993) have been detected in the recent EUV all-sky surveys. EUV detection of these hot WDs depends primarily on their temperature, distance, and the intervening Galactic ISM. Our sample of EUV WDs (whose selection we define below) offers excellent flux contrast in the IR relative to optical; cool companions will almost always be brighter in the K band than the hot WDs.

To know what K mag to expect for the WDs, we benefit from constraints on $\log g$, radius, and T_{eff} derivable from optical spectra for the WDs in our sample, using NLTE model atmosphere fits (Napiwotzki et al. 1993, Napiwotzki 1997). The resulting predictions for K magnitudes allow a direct search for any IR excess from a cool companion. In some cases, IR colors will also provide a preliminary spectral type. Results from the IR survey will be presented in an upcoming paper.

An additional motivation is the study of the white dwarf mass distribution. Since the pioneering work of Koester, Schulz & Weidemann (1979; KSW) it is well established that the masses of white dwarfs cluster in a narrow range around $0.6M_{\odot}$, remarkable given that white dwarfs stem from progenitors with masses ranging from below $1M_{\odot}$ up to $\approx 8M_{\odot}$. Precise knowledge of the white dwarf mass distribution puts constraints on the theory of stellar evolution, especially the poorly understood mass loss process during the final stages of stellar evolution. With two recent exceptions (Beauchamp et al. 1996 and Dreizler & Werner 1996, who analyzed samples of helium-rich DB and DO white dwarfs, respectively) the mass distribution has only been determined for hydrogen-rich DA white dwarfs. However, this is not a severe limitation, because this spectral class comprises about 80% of all known white dwarfs.

The analysis of KSW, along with other follow-up investigations in the early eighties, used photometric data of which the Strömgren and Greenstein multichannel colors were the most suitable. Both systems provide temperature- and gravity-sensitive indices. Alternatively, KSW and others used trigonometric parallax measurements to directly calculate the stellar radius. However, the latter method is practical only for a small sample and suffers from considerable measurement uncertainty. Unfortunately, the photometric indices have their highest sensitivity near 10000K . At this temperature DA white dwarfs have a convective atmosphere and the results depend critically on the adopted parameters of mixing length theory.

The situation improved at the beginning of the nineties, when the development of modern, highly efficient detectors made it possible to obtain high-quality spectra of large numbers of white dwarfs and determine the stellar parameters from a fit to the detailed profiles of the Balmer lines.

This method yields sufficient accuracy for white dwarfs hot enough to have a radiative envelope. The first comprehensive sample of white dwarfs analyzed by this method was presented by Bergeron, Saffer & Liebert (1992; hereafter BSL). As attributed to higher precision of spectroscopic methods, this investigation yielded a white dwarf mass distribution even narrower than found by KSW and other previous studies.

At the same time, Kidder (1991) analyzed a sample of hot DA white dwarfs discovered through positional coincidences of catalogued hot DA white dwarfs in existing soft X-ray databases. Three soft x-ray sources corresponding to white dwarfs were found, having relatively low effective temperatures, $\approx 25,000\text{K}$, determined independently using complementary optical and UV spectroscopy. Kidder et al. (1992) analyzed an expanded sample to derive photospheric He abundances for the hotter objects and establish an effective observational low-temperature threshold for the detection of pure hydrogen DA white dwarfs at soft X-ray wavelengths.

In 1997 three groups (Marsh et al. 1997, M97; Vennes et al. 1997, V97; Finley, Koester & Basri 1997, FKB) published results on the mass distribution of Extreme Ultraviolet (EUV) selected white dwarfs. Due to the selection criterion, these samples contain the hottest white dwarfs ($T_{\text{eff}} > 25000\text{K}$), as cooler white dwarfs do not emit significant EUV radiation. The derived mass distributions in the EUV-selected samples are similar to that of BSL, but show some interesting deviations in detail. The frequency of very high mass white dwarfs is much larger, and that of very low mass white dwarfs much smaller, than in BSL. These findings can at least partly be explained by selection effects (see the discussion in FKB). More serious is a trend of the peak mass with temperature. V97 found that their mass distribution peaks at $0.598M_{\odot}$, while the BSL distribution peaks at $0.568M_{\odot}$, with masses computed using Wood's (1995) mass-radius relation with "thick" layers ($M_{\text{H}} = 10^{-4}M_{\text{WD}}$, $M_{\text{He}} = 10^{-2}M_{\text{WD}}$). This discrepancy diminishes slightly, if the "very thin layer" ($M_{\text{He}} = 10^{-4}M_{\text{WD}}$, no hydrogen layer) mass-radius relations are used (peak masses of $0.556M_{\odot}$ and $0.532M_{\odot}$ for the V97 and BSL sample, respectively). V97 interpreted this as evidence for a very thin hydrogen layer of the DA white dwarfs. However, the effects are small so this result depends strongly on the accuracy of the derived stellar parameters.

FKB estimated the internal accuracy of different analysis methods from Monte Carlo simulations. The precision reachable by Balmer line fitting is very compelling: $\Delta T_{\text{eff}}/T_{\text{eff}} < 0.01$ for $T_{\text{eff}} < 60000\text{K}$. However, for spectra with very high signal-to-noise ratios (S/N), errors introduced by details of the observation and reduction techniques (e.g., extraction, flat fielding, flux and wavelength calibration) might be more important, but are very difficult to determine. Additionally, one has to take into account differences in the model atmosphere calculations and fitting procedure. Together with the results presented in this paper, we now have four samples of hot white dwarfs, analyzed in a similar way, and with significant overlap. This offers the opportunity to determine the real accuracy of the spectral analysis of hot white dwarfs, including many possible systematic effects.

We present the selection criteria of our sample in Sect. 2 and the observations and data reduction procedures in Sect. 3. Details on our model atmospheres are given in Sect. 4. The results and a detailed comparison with the previous analyses of EUV selected white dwarfs are presented in Sect. 5. We finish with a discussion of our results and an outlook.

2. SAMPLE SELECTION CRITERIA

We chose to limit our uniform sample to DAs, for which models provide the best temperature and mass constraints. We start with 73 known DA white dwarfs in the EUVE bright source list (Malina et al. 1994). Excluding sources at low galactic latitudes ($|b| < 15$) and southerly declinations ($\delta < -20$) yields a list of 28 DAs. A similar procedure for non-overlapping DAs listed in the *ROSAT* Wide Field Camera survey Bright Source Catalogue (Pounds et al. 1993) yields 29 objects.

We have removed from our uniform sample 2 well-known stars with published sensitive optical spectrophotometry and IR photometry (Feige 24 and HZ 43). Two DAs with broad line profiles due to magnetic splitting were also excluded (PG 1658+441 and PG 0136+251). Eight known binaries are also excluded from the uniform sample: V471 Tau (Vennes, Christian, & Thorstensen 1998), PG 0824+289, (Heber et al. 1993) HD 74389B (Liebert, Bergeron, & Saffer 1990), RE J1016-052 (V97), PG 1033+464 (GD 123), (Green, Schmidt, & Liebert 1986), RE J1426+500 (V97), RE J1629+780 (Catalan et al. 1995), and IK Peg (Wonnacott et al. 1993), leaving 47 objects.

In this paper, a handful of objects which fell outside the uniform sample definitions just outlined were included for observation. These include the known binaries PG 1033+464 and RE J1629+780 and the magnetic white dwarf PG 1658+441, as well as MCT 0455-2812 which was outside the sample declination limits.

Due to observing constraints (a combination of weather, poor seeing and faint objects, or celestial placement of objects) no spectra were obtained for sample objects RE J0443-034, RE J0916-194, or PG 1040+451. PG 1234+482 was originally classified as a sdB star and thus excluded; it has since been reclassified as a DA (Jordan et al. 1991). The final sample we analyze here thus includes 46 DA white dwarf stars for which we present new model NLTE fits to optical spectra.

We note that since the sample selection was performed, several relevant discoveries pertaining to sample objects have been made. RE J0134-160 (GD984) has central Balmer emission components produced by a dMe companion. RE J1440+750 turns out to be a magnetic DA (Dreizler et al. 1994). The uniform sample will be discussed in a followup paper treating the IR photometry and binary fraction.

3. OBSERVATIONS

Dates are listed for all observations in Table 1. On the nights of 04 - 06 January 1996 we obtained spectra at Steward Observatory's Kitt Peak Station using the Bok 2.3-m reflector equipped with the Boller & Chivens Cassegrain spectrograph and UV-flooded Loral 800×1200 CCD. Most

spectra were dispersed with a 600 l/mm first-order grating used behind a $4.5'' \times 4'$ long slit. The instrumentation provided wavelength coverage $\lambda\lambda 3400\text{--}5600$ at a spectral resolution of $\sim 5\text{\AA}$ FWHM. On the last night of the observing run we employed a new 400 l/mm grating providing wavelength coverage $\lambda\lambda 3500\text{--}6790$ at a spectral resolution of $\sim 7\text{\AA}$ FWHM.

We also obtained spectra at the MMT on Mt Hopkins, 09 - 11 May 1996, using the 300 l/mm grating in first order on the Loral 3k×1k CCD of the Blue Channel spectrograph. This yields coverage from about 3350 to 8800 $\text{\AA}$, and the $2''$ slit width we used resulted in a spectral resolution of about 4 $\text{\AA}$ FWHM. Several objects were kindly obtained for us using the identical instrumental configuration at the MMT by Perry Berlind on 08 April 1997.

Exposures at both telescopes ranged from one to thirty minutes for program stars, and for all observations the long slit was rotated to the parallactic angle according to the calculations of Filippenko (1982). The airmasses were held below 1.5 in almost all cases. All spectra were extracted from the two-dimensional images and reduced to linear wavelength and intensity scales using standard reduction packages in the Image Reduction and Analysis Facility (IRAF). These operations included bias subtraction, flat field division by images obtained by exposing on dome or internal quartz lamps, centroiding and summation of the stellar traces on the two-dimensional images, sky subtraction, wavelength calibration using spectra of He/Ar arc lamps, and absolute flux calibration using the flux standards of Massey et al. (1988). Details of these reduction procedures are given by BSL.

We present in Figs. 1 and 2 our collection of white dwarf spectra. The spectra obtained at the Steward observatory 2.3 m telescope are found in Fig. 1, and the MMT spectra in Fig. 2. Some stars observed repeatedly appear in both figures. The signal-to-noise ratio of our spectra ranges from 35 to 200 with an average of 90. Ninety percent of our spectra have a S/N of at least 45.

4. MODEL ATMOSPHERES

We calculate hydrogen model atmospheres with the NLTE code developed by Werner (1986). Basic assumptions are those of static, plane-parallel atmospheres in hydrostatic and radiative equilibrium. In contrast to the atmospheres commonly used to analyze DA white dwarfs, we relax the assumption of local thermal equilibrium (LTE) and solve the detailed statistical equilibrium instead. As described in Werner (1986), the accelerated lambda iteration (ALI) method is used to solve the set of non-linear equations. The impact of NLTE on white dwarf atmospheres is discussed in detail in Napiwotzki (1997).

H I levels and lines are included in NLTE up to $n = 16$. Line blanketing by the Stark broadened hydrogen lines is taken into account consistently. As the hydrogen atmospheres of DA white dwarfs are stable for $T_{\text{eff}} > 15000\text{ K}$, convection is not included in our atmospheric models. Pressure dissolution of the higher levels is described by the Hummer & Mihalas (1988) occupation probability formalism following the NLTE implementation by Hubeny et al. (1994). The synthetic spectra are computed with the extended VCS broadening tables (Vidal et al. 1970) provided by Schöning & Butler (priv. comm.) and Lemke (1997).

We followed the prescription of Bergeron (1993) and increased the critical ionizing field adopted to calculate the occupation probability by a factor two. The motivation is not a flaw in the Hummer & Mihalas (1988) formalism, but a compensation for the inadequacy of the standard Stark broadening theory when line wings overlap. Our NLTE model grid covers the temperature range $17000\text{ K} < T_{\text{eff}} < 100000\text{ K}$ (stepsize increasing with T_{eff} from 2000 K to 10000 K) and gravity range $6.50 < \log g < 9.75$ (stepsize 0.25).

Although deviations from LTE are small for most DA white dwarfs they become significant for the hottest stars in our sample (cf. Napiwotzki 1997). Since we intend to compare our results with three other samples analyzed by means of LTE atmospheres, we have produced a map with LTE correction vectors. For this purpose we calculated a set of LTE model atmospheres using the technique described in Napiwotzki (1997) of drastically enhancing collisional rates between the atomic levels in the NLTE code. This forces the occupation numbers to be in LTE and guarantees consistency with the NLTE atmospheres.

The synthetic LTE spectra were transformed into “observed” spectra by convolving them with a Gaussian of $5\text{ \AA}$ FWHM, rebinning them to $2\text{ \AA}$ and adding Poisson noise corresponding to a continuum S/N of 100. These simulated spectra were analyzed with the NLTE grid following the procedure outlined in Sect. 5. 1000 simulations were run for every parameter set to eliminate the effect of random errors. The resulting offsets are displayed in Fig. 3. The orientation of the vectors corresponds to the correction, which must be applied to transform LTE results to the NLTE scale.

As expected from the results of Napiwotzki (1997) the differences are negligible for DA white dwarfs cooler than $\approx 40000\text{ K}$, but can be significant for hotter stars. The LTE vectors show the trend that NLTE effects increase with increasing temperature and decreasing gravity (see e.g. Fig. 3 and 4 in Napiwotzki, 1997, for the case of DA white dwarfs). The small corrections found for the models with the highest temperature and lowest gravity seems to contradict this behavior. A look at the line profiles reveals that the NLTE deviations are larger for, say, the $T_{\text{eff}} = 90000\text{ K}$, $\log g = 7.0$ model than the 70000 K , 8.0 model. However, the correction vectors are not a simple function of NLTE deviations measured e.g. as equivalent width difference, but depend also on the way line profiles vary with temperature and gravity. The small corrections found for the high temperature/low gravity models are produced by the cancellation of these effects. We expect a reliable transformation between the LTE and NLTE temperature scales only if the deviations are not too large. A conservative upper limit is $\approx 70000\text{ K}$. A thorough comparison with the previous LTE analyses is presented in Sect. 5.3.

We checked our models by a comparison of our LTE spectra with some DA model spectra kindly provided by D. Koester. Model parameters were $T_{\text{eff}} = 30000\text{ K}$ to 70000 K in 10000 K steps and $\log g = 8.0$ for pure hydrogen models. Input physics are very similar. In particular both model calculations adopt twice the critical ionizing field for the calculation of line profiles. We treated Koester’s model spectra the same way we treated the LTE

spectra above and fitted them with our LTE grid. The result was quite satisfactory: the temperature differences were always below 1.5% and the gravity differences never exceeded 0.03 dex.

Metals were ignored in our calculations, but they can modify the hydrogen line profiles by their effect on the atmospheric structure. Lanz et al. (1996) analyzed the Balmer lines of the hot DA G 191 B2B with pure hydrogen LTE and NLTE atmospheres and an NLTE model with full metal line blanketing. They concluded that the effect of metal line blanketing on the Balmer lines was relatively small, and the difference between LTE and NLTE was found to be the most important effect. A recent study by Barstow et al. (1998), which investigated several hot white dwarfs in the temperature range around 60000 K derived larger metal line blanketing effects of the order of the NLTE effects. Since the LTE analyses of M97, V97, and FKB are based on pure hydrogen models, our results should be consistent with theirs in any case.

5. SPECTRAL ANALYSIS AND RESULTS

Atmospheric parameters of our DA white dwarfs are obtained by simultaneously fitting line profiles of the observed Balmer lines with the NLTE model spectra described above. We use the least-square algorithm described in BSL. The observed and theoretical Balmer line profiles are normalized to a linear continuum (both spectra are F_{λ}) in a consistent manner. Wavelength shifts are determined with a cross-correlation method and applied consistently to each complete spectrum. The synthetic spectra are convolved to the observational resolution with a Gaussian and interpolated to the actual parameters with bicubic splines, and interpolated to the observed wavelength scale.

The atmospheric parameters T_{eff} and $\log g$ are then determined by minimizing the χ^2 value by means of a Levenberg-Marquardt steepest descent algorithm (Press et al. 1986). Several tests revealed that our interpolation routine is rather robust concerning spacing of our model grid and yields reliable results even at the edge of the model grid.

Finally, an estimate of the internal errors can be derived from the covariance matrix. In contrast to BSL, we estimate the noise of the spectra (σ) used for the χ^2 fit from the neighboring continuum of each line. The S/N is adopted to be constant throughout the line.

TABLE 1
ATMOSPHERIC PARAMETERS OF PROGRAM STARS

RE J	Other Names	T_{eff}	$\log g$	$M/M_{\odot}$	Observation Date
0007+331	GD 2	46493±514	7.83±0.05	0.602	6 Jan 1996
0134−160 ^a	GD 984, PHL 1043	44866±667	7.77±0.05	0.572	5 Jan 1996
0237−122	PHL 1400	32077±177	8.45±0.04	0.890	5 Jan 1996
0348−005	GD 50	39508±464	9.07±0.06	1.215	4 Jan 1996
0427+740		48587±1044	7.93±0.08	0.646	5 Jan 1996
0457−280	MCT 0455−2812	51199±786	7.72±0.05	0.570	4 Jan 1996
0512−004		31733±139	7.40±0.03	0.458	5 Jan 1996
0521−102		33186±301	8.60±0.06	0.980	5 Jan 1996
0841+032		38293±252	7.75±0.03	0.544	4 Jan 1996
0902−040		23218±160	7.84±0.02	0.544	4 Jan 1996
0907+505	PG 0904+511	32167±338	8.11±0.07	0.695	4 Jan 1996
0940+502	PG 0937+506	36034±283	7.69±0.04	0.519	9 Apr 1997
0957+852		51311±1348	8.37±0.10	0.866	5 Jan 1996
1019−140		31524±102	7.92±0.02	0.606	9 Apr 1997
1029+450	PG 1026+454	35518±247	7.70±0.04	0.520	9 Apr 1997
1032+532		43587±506	7.95±0.05	0.644	4 Jan 1996
1033−114	G 162−66, LTT 3870	24685±252	7.85±0.03	0.553	4 Jan 1996
1036+460 ^e	GD 123	29361±251	8.02±0.05	0.648	4 Jan 1996
1043+490 ^e		41132±1178	7.94±0.13	0.635	5 Jan 1996
1044+574	PG 1041+580	30338±153	7.81±0.03	0.550	4 Jan 1996
1100+713	PG 1057+719	41104±814	7.84±0.09	0.593	4 Jan 1996
1112+240	Ton 61	39824±636	7.78±0.07	0.563	5 Jan 1996
1122+434	PG 1120+439	26996±151	8.31±0.02	0.803	9 Apr 1997
1126+183 ^e	PG 1123+189	54334±1983	7.76±0.13	0.594	4 Jan 1996
1128−025	PG 1125−026	30699±380	8.24±0.08	0.767	4 Jan 1996
1148+183	PG 1145+188	25758±299	7.91±0.04	0.587	4 Jan 1996
1235+233 ^b	PG 1232+238	46569±523	7.83±0.05	0.602	11+12 May 1996
1257+220 ^b	GD 153	38926±142	7.78±0.02	0.560	4 Jan 1996, 12 May 1996
1336+694	PG 1335+701	29607±87	8.34±0.02	0.824	10 May 1996
1431+370	GD 336	34404±115	7.91±0.02	0.608	11 May 1996
1446+632		37947±254	7.79±0.04	0.562	11 May 1996
1629+780 ^c		41043±338	7.92±0.04	0.627	11 May 1996
1638+350	PG 1636+351	35404±142	7.98±0.02	0.642	10 May 1996
1643+411 ^b	PG 1642+414	28815±81	8.22±0.02	0.753	9 May 1996
1650+403 ^b		38144±211	7.97±0.03	0.643	9 May 1996
1711+664 ^d		48989±757	8.89±0.06	1.141	12 May 1996, 9 Apr 1997
1726+583 ^b	PG 1725+586	53561±542	8.23±0.04	0.795	9 May 1996
1800+683	KUV 18004+6836	44723±424	7.80±0.04	0.585	10 May 1996
1820+580		44099±264	7.78±0.03	0.574	9 May 1996
1845+682	KUV 18453+6819	36120±189	8.23±0.03	0.770	10 May 1996
2116+735 ^b	KUV 21168+7338	50812±354	7.72±0.03	0.569	5 Jan 1996, 9 May 1996
2207+252		26964±174	8.27±0.03	0.779	5 Jan 1996
2312+104	GD 246	53088±968	7.85±0.07	0.624	4 Jan 1996

NOTES: The error estimates were taken from the χ^2 procedure.

^aspectrum slightly contaminated by cool companion; the cores of H β and H γ are excluded from the fit

^bWeighted mean of individual observations

^cred part of spectrum contaminated by cool companion; H β and the core of H γ excluded from fit

^dClose-by optical companion; spectrum apparently not contaminated

^ered part of spectrum contaminated by cool companion; H β excluded from fit

The results are given in Table 1, with illustrative examples shown in Fig. 4. We adopt, for the moment, the usual practice and indicate in Table 1 the internal errors estimated from the quality of the χ^2 fit. However, one should keep in mind that these errors can only serve as lower limits. We will show below (Sect. 5.3) that these formal errors derived from the χ^2 fit significantly underestimate the real errors.

External errors can be estimated from multiple observations and analysis of the same star. We obtained repeat observations for a subsample of 6 stars, for which results are given in Table 2. The gravity values of all six stars agree within the estimated internal errors. The same is true for four temperature comparisons, too. However, the differences found for RE J1650+403 and RE J2116+735 are significantly larger. This is in line with the external errors we estimate from a comparison with the studies of M97, V97, and FKB (see Sect. 5.3).

5.1. Binaries

Five stars in our sample, RE J0134-160, RE J1036+460, RE J1043+490, RE J1126+183, and RE J1629+780, show clear signs of binarity in the red part of their spectra. Two more stars, RE J1711+664 and RE J2207+502, are members of visual binaries.

RE J1629+780: The red part of the spectrum of RE J1629+780 is heavily contaminated by a M type main sequence companion. The composite spectrum and the spectrum of the M star after subtracting the white dwarf component are shown in Fig. 5. The characteristic bands of TiO are easily recognizable. Catalán et al. (1995) determined a spectral type dM4. The H α and H β lines are seen in emission, which indicates a chromospherically active Me star. Sion et al. (1995) detected a flare-like increase of the Balmer line emission.

The comparison in Fig. 5 demonstrates that the blue part ($\lambda < 4400 \text{ \AA}$) of the white dwarf spectrum is not disturbed by the M dwarf. Thus, we excluded H β and the core of H γ from the fit and derived the parameters $T_{\text{eff}} = 41000 \text{ K}$ and $\log g = 7.92$ from H γ to H η . These parameters are in reasonable agreement with the results of Catalán et al. ($T_{\text{eff}} = 41800 \text{ K}$; $\log g = 8.0$), who however included H β , and Kidder (1991; $T_{\text{eff}} = 42500 \text{ K}$, $\log g = 7.6$), who fitted Lyman- α and the Balmer lines H β and H γ .

RE J1036+460, RE J1043+490, RE J1126+183: Three more white dwarfs of our observed sample are known binaries (RE J1036+460, RE J1126+183: Green, Schmidt, & Liebert 1986, RE J1043+490: Schwartz et al. 1995) and show red excesses in our spectra: Since the H β lines of the white dwarfs are contaminated they were excluded from the fit. H γ and higher Balmer lines are virtually uncontaminated. A discussion of these stars and newly discovered binaries will be given in a forthcoming paper.

RE J0134-160 (GD 984): Although our spectrum ends at only $5600 \text{ \AA}$ a red excess is obvious. Subtraction of the theoretical white dwarf flux leaves an M star spectrum. The M dwarf contribution is much smaller than

in RE J1629+780 and we excluded only the cores of H β and H γ , filled in by the Balmer line emission, from the fit. Bues & Aslan (1995) suspected a hot third component in RE J0134-160. However, outside of the H β and H γ cores, the Balmer lines are well reproduced by our best fit without any indication of a third component. Indeed, a subdwarf component as suggested by Bues & Aslan (1995) is almost certainly not present, since its flux would dominate in the blue.

RE J1711+664: Since a late type star is separated from the white dwarf by only $2''.5$, we took care to get an uncontaminated white dwarf spectrum. We obtained a spectrum under good seeing conditions. No excess is present up to the red limit at $8500 \text{ \AA}$.

RE J2207+252: This white dwarf has a red companion $8''.5$ away. Schwartz et al. (1995) estimated spectral type K4V and distance 65 pc from its colors. With the parameters from Table 1 a white dwarf distance of 62 pc results. Thus it is likely that both stars form a physical pair. The angular separation corresponds to $\approx 500 \text{ AU}$.

5.2. Magnetic white dwarfs

Two white dwarfs of our sample show Zeeman splitting of the Balmer line cores, indicative of a magnetic field. RE J1659+440 (PG 1658+441) is a well known star, already analyzed by Schmidt et al. (1992). The discovery of the magnetic nature of RE J1440+750 (HS 1440+7518) was announced by Dreizler et al. (1994; note the naming confusion corrected in Dreizler et al. 1995: HS 1412+6115 should have been HS 1440+7518). Although RE J1440+750 was analyzed by V97, they did not remark on its magnetic nature. This is likely due to the lack of coverage of the H α line, which displays the most pronounced Zeeman effect.

Flux calibrated spectra of the magnetic white dwarfs are displayed in Fig. 6. The PG 1658+441 analysis of Schmidt et al. (1992) resulted in $T_{\text{eff}} = 30500 \text{ K}$ and $\log g = 9.35$. The magnetic splitting was best reproduced by a $3.5 \text{ megagauss (MG)}$ dipole inclined 60° to the line of sight (producing a mean surface field strength $B_S = 2.3 \text{ MG}$). From the linear Zeeman effect we estimated a mean magnetic strength of 8 MG for RE J1440+750, consistent with the estimate given in Dreizler et al. (1994).

The temperature and gravity of RE J1440+750 were derived from a fit of the higher Balmer lines H γ to H η , which are less affected by the magnetic splitting. Results are given in Table 3 supplemented by the Schmidt et al. (1992) fit of PG 1658+441. Our own fit gave similar results albeit with lower accuracy. Our fits of RE J1440+750 can only provide a rough estimate of the stellar parameters. Accurate results can only be expected from a detailed treatment of the magnetic effects.

5.3. Internal, external, and systematic errors

We have now presented the results of a homogeneous analysis of a sample of 46 hot, EUV selected white dwarfs based on Balmer line fitting. Three other large samples analyzed with the same method were recently published by M97, V97, and FKB. Since considerable overlap exists

TABLE 2
RESULTS OF REPEATED OBSERVATIONS

RE	Observation 1		Observation 2	
	T_{eff}	$\log g$	T_{eff}	$\log g$
1235+233	46308±837	7.85±0.08	46737±670	7.82±0.06
1257+220	39349±317	7.76±0.04	38820±159	7.78±0.02
1643+411	28813±120	8.23±0.02	28818±111	8.21±0.02
1650+403	37798±245	7.94±0.04	39126±413	8.01±0.05
1726+583	52712±927	8.27±0.06	54003±669	8.20±0.05
2116+735	50131±384	7.71±0.03	54604±906	7.76±0.06

NOTES: The error estimates were taken from the χ^2 fitting procedure.

TABLE 3
PARAMETERS OF THE MAGNETIC WHITE DWARFS

RE	other names	T_{eff}	$\log g$	$M/M_{\odot}$	B_{S}	Observation Date
1440+750	HS 1440+7518	36154±275	8.87±0.05	1.128	7.7	10 May 1996
1659+440 ^a	PG 1658+441	30510±200	9.36±0.07	1.311	2.3	10 May 1996

^aParameters from Schmidt et al. 1992

between all four samples, this allows a direct check for systematic errors and the individual scatter on a star by star basis for white dwarfs hotter than 25000 K.

Since in contrast to previous works, our analysis is based on NLTE model atmospheres, we applied the correction vectors given in Fig. 3 to correct for the LTE assumption. Since for the hottest white dwarfs these corrections become large, while the accuracy of temperature and gravity estimates decreases for both LTE and NLTE analyses, one should exclude comparison of stars with $T_{\text{eff}} > 70000$ K. This does not affect our sample, which has a maximum temperature closer to 54000 K.

Differences (after correction to NLTE) in T_{eff} and $\log g$ between studies for stars in common with M97, V97, and FKB are displayed in Fig. 8 as function of T_{eff} . The magnetic white dwarfs and the binaries, which show significant contamination of the white dwarf spectrum by the companion, are excluded. One can now focus on systematic differences between pairs of studies, e.g. our results (NGS) versus FKB (FKB–NGS), V97–NGS, M97–NGS, FKB–M97, M97–V97, and so on. However, we chose another approach and performed an optimization that simultaneously took into account all values from all stars in common between any samples. In other words, the values given for the systematic differences between our study and the M97, V97, and FKB samples form a system for direct transformation between, say, FKB and the three other samples. The running averages computed this way are plotted in Fig. 8 as a solid line. The actual value was computed in 1000 K steps for all white dwarfs in the respective temperature intervals $T_{\text{eff}} \pm 10\%$. Although this curve does not represent the best fit to the data plotted in Fig. 8 alone, it is a fairly good representation of the differences computed directly between our measurements and those of M97, V97, and FKB.

Since the distribution is highly non-Gaussian with many outliers, as evident in Fig. 8, we decided to adopt an underlying Lorentzian (or Cauchy) distribution for the optimization. The tails of the Cauchy distribution are much larger than that of the corresponding Gaussian, yielding a much lower weight for deviant points (see discussion in Press et al. 1986). The dotted lines represent the 1σ confidence interval of the mean, computed conservatively from the rms deviations.

First, we notice a considerable scatter, larger than expected from the internal error estimates (see the discussion below). If one ignores the hot end, the agreement between the FKB and our temperature scale is good; differences are below 1%, smaller than the maximum model differences to the Koester models (cf. Sect. 4), which were used by FKB. The same atmospheres are used in M97, and it is therefore surprising that significant differences with M97 are present. These trends are most likely caused by different reduction and analysis techniques. Offsets of the same order are found in our comparison with V97, where a different LTE model atmosphere code is used. Although basically the same input physics is included, this might at least partly explain those shifts in T_{eff} and $\log g$.

Our results are quantified in Tables 4 and 5. We divided the white dwarfs into three groups according to their temperature: a cool group with $T_{\text{eff}} < 30000$ K, a hotter one with $30000 \text{ K} < T_{\text{eff}} < 45000$ K, and the hottest consid-

ered group with $45000 \text{ K} < T_{\text{eff}} < 70000$ K. NLTE effects are still negligible in the range of effective temperatures of the two coolest groups. Mean shifts and the confidence range of the mean (computed as described above) are provided for these groups and the complete sample (except stars with $T_{\text{eff}} > 70000$ K). The shifts discussed above and shown in Fig. 8 are statistically significant and reach values of $\approx 5\%$ in T_{eff} in the hottest bin. In $\log g$ the difference between our results and M97 reach ≈ 0.1 dex for the coolest bin. However, we emphasize that all four analyses are based on state-of-the-art model atmospheres and χ^2 fitting techniques. Since there are no strong arguments to favor or discard one analysis, it seems these are the systematic shifts characteristic of modern analyses of hot white dwarfs.

If we take the systematic shifts into account, we can use the samples to derive estimates of the observational scatter, which can be compared with the internal error estimates. Since we have approximately the same scatter for different combinations of the samples, we compute a mean scatter σ_{diff} for all possible combinations (weighted by the number of stars in common). For this purpose we correct for the systematic shifts calculated for each of the three T_{eff} bins, whose temperature intervals are given in Table 6. The individual measurement errors σ_{ind} add quadratically, and if we assume that the inherent scatter is the same for all analyses (and we found no reason to discard this assumption), the individual measurement errors can be estimated as $\sigma_{\text{ind}} = \frac{1}{\sqrt{2}}\sigma_{\text{diff}}$. Not surprisingly, the smallest scatter is found for the “cool” group ($T_{\text{eff}} < 30000$ K) with $\sigma(T_{\text{eff}}) = 2.3\%$ and $\sigma(\log g) = 0.07$ dex. It increases to $\sigma(T_{\text{eff}}) = 3.3\%$ and $\sigma(\log g) = 0.13$ dex for the hottest bin. This trend is expected from the statistical analysis presented in FKB (their Fig. 1). However, the values are larger by a factor of three or more than the internal parameter errors for a well exposed spectrum (see e.g. Table 1). Therefore, we conclude that the accuracy is not limited by the noise for good spectra, and we suggest that other effects, such as details of the extraction or fluxing and normalization procedures, contribute more. Considering these systematic uncertainties, the 0.3 dex difference between the gravity determinations of FKB and Napiwotzki et al. (1993) for HZ 43 A is only a 1.5σ deviation and therefore not as serious as considered by FKB.

5.4. Mass distribution

5.4.1. Derivation and Sample Comparisons

Once the temperature and gravity of the white dwarfs are known the mass can be determined from theoretical mass-radius relations. The recent investigations of M97, V97, and FKB based their interpretation on the model sequences of Wood (1995). The applied models have H- and He-layer masses of $10^{-4}M_{\text{WD}}$ and $10^{-2}M_{\text{WD}}$, respectively (“thick layers”) and a carbon core. These model sequences are based on a set of pre-white dwarf models with end masses of $0.6M_{\odot}$, $0.8M_{\odot}$ and $0.95M_{\odot}$ computed by Kawaler (see Wood 1994). Starting models for other masses were constructed by homology transformations.

The mass-radius relations we use are based on the evolutionary calculations of Blöcker (1995). These models are calculated *ab initio* from the main sequence. The

TABLE 4
SYSTEMATIC TEMPERATURE DIFFERENCES

	M97-NGS $\Delta T_{\text{eff}}/T_{\text{eff}}$	V97-NGS $\Delta T_{\text{eff}}/T_{\text{eff}}$	FKB-NGS $\Delta T_{\text{eff}}/T_{\text{eff}}$
$T_{\text{eff}} < 30000 \text{ K}$	-0.023 ± 0.006	0.032 ± 0.005	0.007 ± 0.005
$30000 < T_{\text{eff}} < 45000 \text{ K}$	-0.013 ± 0.003	0.017 ± 0.003	0.004 ± 0.003
$45000 < T_{\text{eff}} < 70000 \text{ K}$	0.036 ± 0.006	0.050 ± 0.006	0.046 ± 0.007
all with $T_{\text{eff}} < 70000 \text{ K}$	-0.006 ± 0.003	0.024 ± 0.003	0.012 ± 0.003

NOTES: For each temperature range, each column gives the average difference over the respective interval, and the confidence interval of the average difference.

TABLE 5
SYSTEMATIC GRAVITY DIFFERENCES

	M97-NGS $\Delta \log g$	V97-NGS $\Delta \log g$	FKB-NGS $\Delta \log g$
$T_{\text{eff}} < 30000 \text{ K}$	-0.112 ± 0.017	-0.001 ± 0.016	-0.064 ± 0.018
$30000 < T_{\text{eff}} < 45000 \text{ K}$	-0.086 ± 0.017	0.026 ± 0.016	-0.034 ± 0.016
$45000 < T_{\text{eff}} < 70000 \text{ K}$	-0.066 ± 0.030	-0.067 ± 0.026	-0.062 ± 0.027
all with $T_{\text{eff}} < 70000 \text{ K}$	-0.084 ± 0.013	-0.002 ± 0.012	-0.047 ± 0.012

NOTES: For details see Table 4

TABLE 6
MEAN SCATTER σ_{ind} OF TEMPERATURE AND GRAVITY DETERMINATION

	$\sigma(T_{\text{eff}})$	$\sigma(\log g)$
$T_{\text{eff}} < 30000 \text{ K}$	0.023	0.075
$30000 < T_{\text{eff}} < 45000 \text{ K}$	0.023	0.101
$45000 < T_{\text{eff}} < 70000 \text{ K}$	0.034	0.128
all with $T_{\text{eff}} < 70000 \text{ K}$	0.026	0.107

resulting white dwarfs show important differences compared to Wood’s models. In contrast with the carbon core models widely used for mass determination, the degenerate core contains a mass-dependent mixture of carbon and oxygen (and trace elements). However, note that Wood (1995) also presented some white dwarf sequences with C/O cores. The H- and He-layer masses resulting from evolutionary calculations depend on the stellar mass (Blöcker 1995; Blöcker et al. 1997). While the canonical thick hydrogen layer mass of $10^{-4}M_{\odot}$ is met for a $0.6M_{\odot}$ white dwarf, it decreases from several $10^{-3}M_{\odot}$ for $0.3M_{\odot}$ down to $10^{-6}M_{\odot}$ for $1.0M_{\odot}$. This results in a mass-radius relation which is steeper than a relation based on a constant layer mass. Consequently, the derived mass distribution will be narrower, if evolutionary layer masses are used.

The lower mass limit for a C/O core white dwarf is $0.46M_{\odot}$, the limiting mass for central helium burning in low mass stars (Sweigart et al. 1990). Thus, white dwarfs with a lower mass possess a helium core. At the current age of the universe He white dwarfs have not been produced by single star evolution, but are the result of binary evolution where the hydrogen-rich envelope was stripped away along the (first) red giant branch (Kippenhahn 1967; Iben & Tutukov 1986). We use the recent evolutionary models of He white dwarfs calculated by Driebe et al. (1998) based on this scenario.

The combined tracks of Blöcker (1995) and Driebe et al. (1998) cover the mass range $0.18M_{\odot}$ to $0.94M_{\odot}$. We supplemented this set with the 1.0, 1.1, and $1.2M_{\odot}$ carbon core sequences of Wood (1995) with “thin” layers. Since the hydrogen envelope mass decreases with increasing white dwarf mass in Blöcker’s models, their highest mass models effectively correspond to “thin layer” models. Since high mass white dwarfs are already close to the zero-temperature configuration, the remaining departures from consistent evolutionary calculations can be ignored for practical purposes.

The position of the analyzed white dwarfs in the temperature/gravity plane is shown in Fig. 9 along with the tracks used for the mass determination. The individual white dwarf masses are given in Table 1, and the resulting mass distribution is shown in Fig. 10. We have also redetermined masses for the BSL sample with the Blöcker/Driebe mass-radius relations and supplement Fig. 10 with the resulting distribution.

Our mass distribution possesses the same basic features as found in the EUV selected samples of M97, V97, and FKB. The sharp peak centered at $\approx 0.59M_{\odot}$ is in principal agreement with the earlier investigations of KSW, Weidemann & Koester (1984), and BSL, with a sharp falloff towards lower masses, and a less steep decline towards higher masses with a tail of high mass white dwarfs. Considering the relatively large observational scatter we found in Sect. 5.3, the *underlying* distribution may be extremely sharp. The mean mass is $0.67M_{\odot}$. However, this value is strongly biased by the few very high mass white dwarfs.

We decided to follow the recipe of FKB and fitted the mass peak with a Gaussian. Although a Gaussian is not a good representation of the white dwarf mass distribution, it gives a robust estimate of the peak mass. The fit result is, in principle, dependent on the adopted binning

of the mass intervals. However, in some test calculations we found effects exceeding a few $0.001M_{\odot}$ only if the bin width was larger than $0.05M_{\odot}$. The values given in this paper were derived by fitting Gaussians to virtually unbinned mass distributions (formal bin width $0.001M_{\odot}$).

We were concerned by the possibility that the different fraction of low and high mass white dwarfs found in different samples (see discussion below) may skew the Gaussian fit of the main peak to higher or lower masses. We tested this by fitting the BSL, and FKB and our mass distributions with multi-component Gaussians, which fitted the secondary high and low mass peaks separately. We found no deviation higher than $0.003M_{\odot}$ in any case and concluded that the single Gaussian fit of the main peak yields a rather stable estimate.

With this method, we derived a peak mass of our sample of $0.589M_{\odot}$. We reanalysed the FKB, M97, and V97 samples with our mass-radius relations. We applied the corrections to NLTE and excluded the white dwarfs with temperatures in excess of 70000 K. The peak masses are $0.555M_{\odot}$ (FKB), $0.535M_{\odot}$ (M97), and 0.582 (V97). These differences reflect the systematic differences in T_{eff} and $\log g$ determinations discussed above and probably different selection criteria in the case of FKB.

The probable He core white dwarf RE J0512-004 is the only object in our sample with a mass below $0.5M_{\odot}$, but four white dwarfs have masses in excess of $1.0M_{\odot}$. The frequency of low and high mass white dwarfs found in our and other EUV selected samples is qualitatively different from that found in optically selected samples such as BSL. BSL detected in their sample of 129 white dwarfs 16 objects with masses below the limiting mass for C/O core white dwarfs and a total of 28 white dwarfs with $M < 0.5M_{\odot}$. These numbers are reduced to 10 and 16 respectively, if we redetermine the masses with the mass-radius relations of Blöcker (1995) and Driebe et al. (1998). However, that is still a much higher fraction than we find in our EUV selected sample.

5.4.2. Cooling Rates and Detection Probability

Why do optical and EUV selected samples contain different fractions of high and low mass white dwarfs? All helium white dwarfs from the BSL sample have temperatures below 30000 K; the only helium white dwarf candidate in our sample has $T_{\text{eff}} \approx 32000$ K. This implies that the detection probability of low mass white dwarfs in EUV selected samples is much lower, because the fraction of low mass white dwarfs at higher T_{eff} is much lower than it is for $T_{\text{eff}} < 30000$ K. This viewpoint is supported by the helium white dwarf sequences calculated by Driebe et al. (1998). We have plotted the variation of temperatures with time for white dwarfs with helium and C/O cores in Fig. 11. The detection probability in a given evolutionary stage goes as the inverse of the rate of temperature decrease at that stage. New He white dwarfs cool down rapidly to a certain (mass dependent) temperature, whereafter the cooling rate drops drastically and the star remains on a temperature plateau for a long time. This behavior is most pronounced for the lowest masses. Hydrogen shell burning plays an important role for this plateau phase. Although the energy production by hydrogen burning drops down dramatically after the star enters the white dwarf

sequence, it still produces a significant fraction of the total luminosity, enough to bring the cooling to a standstill. We can conclude from Fig. 11 that it is extremely unlikely to detect a helium white dwarf during its first rapid cooling phase. It is easier to find them during their plateau phase or afterwards. This means there is a strong bias towards detecting helium white dwarfs in samples of relatively “cool” DA stars (like BSL) instead of the overall much hotter EUV selected samples.

We can explain in a similar fashion why high mass white dwarfs are found preferentially in the $30000\text{ K} < T_{\text{eff}} < 50000\text{ K}$ range. Due to the drop in neutrino cooling, the evolutionary speeds of massive white dwarfs are relatively low in this range, as demonstrated by the $0.940M_{\odot}$ track, the most massive stellar remnant calculated by Blöcker (1995). This increases the detection probability of more massive white dwarfs in EUV selected samples.

5.4.3. Interstellar Absorption

Even if we restrict our comparisons to hot subsamples of white dwarfs, the fraction of massive white dwarfs is much higher in the EUV selected samples than it is in optically selected ones. An extensive discussion of this problem is carried out by FKB. FKB argue that the number of high mass white dwarfs discovered per sky area is similar for the whole sky EUV surveys and the optical Palomar-Green (PG) survey (Green et al. 1986), which covered roughly 25% of the sky. In other words, a whole sky version of the PG survey should have discovered all but one of the EUV detected white dwarfs with $M > 1.1M_{\odot}$. At a given temperature, white dwarfs with lower masses are larger, consequently more luminous, and detectable to larger distances. For instance at 30000K, a white dwarf of $0.5M_{\odot}$ is detectable at distances $\sim 30\%$ larger than a $0.7M_{\odot}$ white dwarf. At the maximum distance sampled by, e.g., the PG survey, the probability is very high that interstellar matter effectively absorbs all EUV radiation. Thus the dominant effect in EUV surveys is not a selection for massive white dwarfs as suggested by V97, but a selection *against* low mass white dwarfs, due to a sample volume strongly affected by interstellar absorption.

5.4.4. Temperature Dependence of the Derived Mass Peak

FKB noted a moderate apparent trend of the white dwarf peak mass with temperature. Since their sample also included optically selected white dwarfs, they analyzed a considerably higher fraction of stars with $T_{\text{eff}} < 25000\text{ K}$ and therefore have a larger temperature baseline than pure EUV samples. Analogous to this finding, V97 found an $0.03M_{\odot}$ offset between the peak of the mass distribution in their EUV selected sample and the (cooler) white dwarfs analyzed by BSL. In both cases this temperature dependence decreases if Wood’s models with “very thin” layers ($M_{\text{He}} = 10^{-4}M_{\text{WD}}$, no hydrogen layer) instead of the canonical thick layer models ($M_{\text{He}} = 10^{-2}M_{\text{WD}}$, $M_{\text{H}} = 10^{-4}M_{\text{WD}}$) are used. However, part of the trend noted in V97 may be related to the systematic differences between the analyses discussed in Sect 5.3. Since V97 compared results from two different analyses, it is not possible to judge if the offset is

real or just an artifact. The case of FKB is stronger, because they based their case on a homogeneously analyzed sample. FKB warned that the temperature dependence of the mass distribution peak may be caused by inadequacies remaining in the model atmospheres. A more detailed discussion is given in the next section.

Some evidence that part of the temperature trend stems from the analyses comes from our intercomparisons presented in Sect 5.3. We showed that systematic differences between the four investigated samples exist that vary with effective temperature. This might mimic a temperature dependence of the sample peak mass. A gravity offset of 0.1 dex transforms into mass offsets of $0.050M_{\odot}$, $0.044M_{\odot}$, and $0.034M_{\odot}$ for a $0.6M_{\odot}$ white dwarf with 25000 K, 40000 K, and 60000 K, respectively. Even a relatively small systematic $\log g$ difference of 0.05 dex (cf. Table 5) corresponds to a $0.02M_{\odot}$ offset. The scatter σ of *individual* gravity determinations reported in Table 6 corresponds to $\sigma(M) \approx 0.04M_{\odot}$, nearly independent of T_{eff} .

5.4.5. Cooling Tracks

Another part of the T_{eff} dependence of the mass peak may be caused by the use of Wood’s cooling tracks by FKB and V97. These tracks are not completely self-consistent since not all model sequences were not calculated *ab initio* from the main sequence. This is not important for cooler white dwarfs but may cause deviations from *ab initio* tracks especially for the hottest white dwarfs, where structure depends sensitively on the evolutionary history (Blöcker & Schönberner 1990).

We redetermined the masses of the BSL white dwarfs with the Blöcker/Driebe mass-radius relations and derived a peak mass of $0.559M_{\odot}$. That is $0.03M_{\odot}$ lower than our peak mass, but having in mind the range of mass determination derived for the EUVE selected samples, we cannot consider this a significant difference. FKB divided their sample into cool ($T_{\text{eff}} < 35000\text{ K}$) and a hot ($35000\text{ K} < T_{\text{eff}} < 75000\text{ K}$) subsamples and derived a $0.029M_{\odot}$ higher peak mass for the hot sample. Our reanalysis of the FKB LTE results with the Blöcker/Driebe mass-radius relation yields virtually the same offset: $0.025M_{\odot}$. However, the difference is brought down to $0.010M_{\odot}$ ($0.550M_{\odot}$ vs. $0.560M_{\odot}$) if we apply corrections to NLTE. One can imagine that a difference of this order (if significant at all) can easily be produced by our neglect of metallicity effects (cf. Lanz et al. 1996, Barstow et al. 1998).

Therefore, given the combination of sample selection and systematic effects in analyses to date, our results do not confirm the presence of intrinsic, systematic mass differences between hot and cool white dwarfs.

6. CONCLUSIONS

We have obtained temperatures, gravities, and masses for a sample of 46 extreme ultraviolet selected DA white dwarfs. These data complement a near infrared survey for low mass companions. The stellar parameters were determined by fitting the hydrogen Balmer line profiles with NLTE model spectra. A map of LTE correction vectors was constructed, which allows the transformation of LTE results to the NLTE scale. These shifts become impor-

tant for temperatures above 50000 K (see also Napiwotzki 1997).

Three recent analyses of DA white dwarfs have a considerable overlap with our sample and allow a direct check for systematic errors and individual measurement scatter. Although all four analyses applied similar model atmospheres and fitting techniques, we recognized systematic shifts up to 5% in temperature and 0.1 dex in gravity determinations. These systematic errors have to be considered when the results are interpreted and compared to other studies. The individual measurement errors increase with temperature: $\sigma(T_{\text{eff}})$ from 2.3% to 3.3% and $\sigma(\log g)$ from 0.07 dex to 0.13 dex. This is larger than the typical errors computed with χ^2 fit procedures for a well-exposed spectrum and indicates that accuracy is limited by noise in combination with other effects such as details of the extraction, fluxing or normalization procedures. It is unclear if these σ values indicate lower limits, or scale (at least partially) with the S/N ratio of the spectra. It will be a challenge to resolve this question. Repeated observations performed by one observer, at one telescope, reduced the same way and analysed with one method as performed by BSL, FKB, and us (Table 2) do not reveal all effects. Independent observations analysed independently are necessary.

Masses have been inferred from theoretical mass-radius relations based on the evolutionary calculations of Blöcker (1995) for C/O white dwarfs and Driebe et al. (1998) for He white dwarfs. An important feature of these calculations are the hydrogen and helium layer masses which depend on stellar mass. We find a sharp peak centered at $\approx 0.59 M_{\odot}$ in agreement with the previous investigations of KSW and BSL. We redetermined masses of the BSL white dwarfs with the Blöcker/Driebe mass-radius relation and derived a peak mass of $0.56 M_{\odot}$. At first glance this seems to confirm the systematic offset reported by V97. However, this offset could be explained by system-

atic differences of the model atmosphere analyses as well. Our reanalysis of the homogeneous FKB sample showed that the temperature dependence of the mass peak nearly vanishes, when corrected for their LTE assumption. We conclude that the observational data presented here and in similar studies are well explained by canonical stellar evolution theory, i.e. white dwarfs with thick envelopes.

We find only one object (RE J0512-004) with a mass below $0.5 M_{\odot}$, which is a possible helium core white dwarf, but four white dwarfs with masses in excess of $1.0 M_{\odot}$. This ratio of high- and low-mass white dwarfs is quite different from that found by BSL. This can partly be understood as the result of variable evolutionary time scales of high and low mass white dwarfs. We agree with FKB that another part can be explained by a selection *against* low mass white dwarfs in EUV selected samples.

The principal aim of our project is the search for and study of cool main sequence companions of our white dwarf sample. Five white dwarfs already show red contamination in our optical spectra. A comprehensive discussion of these and other binaries will be given in a forthcoming paper presenting IR photometry for the white dwarf sample. In this article we provided the basic white dwarf data necessary to interpret our binary sample.

We thank T. Driebe, F. Herwig, and T. Blöcker for providing us with their tracks and computing the white dwarf masses and D. Koester, who made some model spectra available for our model comparison. We are grateful to Steward Observatory for an unexpected, generous award of six nights of unclaimed 2.3-m time, which made possible simultaneous optical and infrared observations of our sample of stars. We thank Perry Berlind for obtaining several white dwarf spectra. PJG acknowledges support through NASA Contract NAS8-39073 (ASC).

REFERENCES

- Barstow M. A., Hubeny I., Holberg J. B. 1998, MNRAS, 299, 520
 Beauchamp A., Wesemael F., Bergeron P., Liebert J., Saffer R. A. 1996, in: Hydrogen-deficient stars, eds. C. S. Jeffery & U. Heber, ASP Conf. Ser. Vol. 96, p. 295
 Bergeron P. 1993, in: White dwarfs: advances in observation and theory, ed. M. A. Barstow, Kluwer, Dordrecht, p. 267
 Bergeron P., Saffer R. A., & Liebert J. 1992 ApJ, 394, 228 (BSL)
 Blöcker T. 1995 A&A, 299, 755
 Blöcker T. & Schönberner D. 1990, A&A, 240, L11
 Blöcker T., Herwig F., Driebe T., Bramkamp H., & Schönberner D. 1997, in: White dwarfs, eds. I. Isern et al., Kluwer, Dordrecht, p. 57
 Bues I., Aslan T. 1995, in: White dwarfs, eds. D. Koester & K. Werner, Springer, Berlin, p. 259
 Burleigh, M. R., Barstow, M. A., & Fleming, T. A. 1997, MNRAS, 287, 381
 Catalán M. S., Sarna M. J., Jomaron C. M., & Smith R. C. 1995, MNRAS, 275, 153
 deKool, M. & Ritter, H. 1993, A&A, 267, 397
 Dreizler S., Heber U., Jordan S., & Engels D. 1994, in: Hot stars in the galactic halo, eds. S. J. Adelman, A. R. Upgren, C. J. Adelman, Cambridge University Press, p. 228
 Dreizler S. & Werner K. 1996 A&A, 314, 217
 Dreizler S., Werner K., & Heber U. 1995, in: White dwarfs, eds. D. Koester & K. Werner, Springer, Berlin, p. 160
 Driebe T., Schönberner D., Blöcker T., Herwig F. 1998, A&A, 339, 123
 Filippenko A. V. 1982, PASP, 94, 715
 Finley, D. S., et al. 1993, ApJ, 417, 259
 Finley D. S., Koester D., & Basri G. 1997 ApJ 488, 375 (FKB)
 Green, P. J. & Margon, B. 1994, ApJ, 423, 723
 Green R. F., Schmidt M., & Liebert J. 1986 ApJS 61, 305
 Hubeny I., Hummer D. G., & Lanz T. 1994, A&A, 282, 151
 Hummer D. G. & Mihalas D. 1988 ApJ, 331, 794
 Iben, I. Jr. & Tutukov A. V. 1986 ApJ, 311, 742
 Jordan S., Heber U., Weidemann V. 1991, in: White dwarfs, ed. G. Vauclair & E. M. Sion, Kluwer, Dordrecht, p. 121
 Kidder, K. M. 1991, PhD thesis, University of Arizona
 Kidder, K. M., Holberg, J. B., Barstow, M. A., Tweedy, R. W., & Wesemael, F. 1992, ApJ, 394, 288
 Kippenhahn R., Kohl K., & Weigert A. 1967, Zeitschrift f. Astrophys. 66, 58
 Koester D., Schulz H., & Weidemann V. 1979, A&A, 76, 262 (KSW)
 Lanz T., Barstow M. A., Hubeny I., & Holberg J. B. 1996, ApJ, 473, 1089
 Lemke M. 1997 A&AS 122, 285
 Liebert J., Bergeron P., & Saffer R. A. 1990 PASP, 102, 1126
 Malina, R. et al. 1994, AJ, 107, 751
 Marsh M. C., Barstow, M. A., Buckley D. A. et al. 1997, MNRAS, 286, 369 (M97)
 Massey P., Strobel S., Barnes J. V., & Anderson E. 1988, ApJ, 328, 315
 Napiwotzki R. 1997 A&A, 322, 256
 Napiwotzki R., Barstow M. A., Fleming T., Holweger H., Jordan S., & Werner K. 1993 A&A, 278, 478
 Pounds, K. et al. 1993, MNRAS, 260, 77
 Press W. H., Flannery B. P. Teukolsky S. A., & Vetterling W. T. 1986, Numerical Recipes, Cambridge University Press
 Schmidt G. D., Bergeron P., Liebert J., & Saffer R. A. 1992, ApJ, 394, 603
 Schwartz R. D., Dawkins D., Findley D., & Chen D. 1995 PASP, 107, 667
 Sion E. M., Holberg J. B., Barstow M. A., & Kidder, K. M. 1995, PASP, 107, 232

- Sweigart A. V., Greggio L., & Renzini A. 1990 ApJ, 364, 527
Vennes, S. & Thorstensen, J. R. 1994, ApJ, 433, L29
Vennes S., Thejll P. A., Galvan G. G., & Dupuis J. 1997, ApJ, 480, 714 (V97)
Vennes S., Christian D. J. & Thorstensen J. R. 1998 ApJ, 502, 763
Vidal C. R., Cooper J., & Smith E. W. 1970 JQSRT 10, 1011
Weidemann V. & Koester D. 1984, A&A, 132, 195
Werner K. 1986 A&A, 161, 177
Wonnacott D., Kellett B. J., & Stickland D. J. 1993, MNRAS, 262, 277
Wood, M. 1994, in IAU Coll. 147: The Equation of State in Astrophysics, eds. G. Chabrier and E. Schatzmann (Cambridge U. Press: Cambridge) p. 612.
Wood M. 1995, in: White dwarfs, eds. D. Koester & K. Werner, Springer, Berlin, p. 41

Fig. 1.— Spectra of the white dwarfs observed at Steward observatory

Fig. 2.— Spectra of the MMT sample

Fig. 3.— LTE offsets. The differences are magnified three times. The vectors give the correction which must be applied to transform LTE results to the NLTE scale.

Fig. 4.— Balmer line fits for a representative set of white dwarfs.

Fig. 5.— Flux calibrated spectrum of the DA+M pairs RE J0134-160, RE 1036+460, RE J1043+490, RE J1126+183, and RE J1629+780. From top to bottom we show the composite spectrum, the best fit of the white dwarf, and the resulting spectrum of the M star. The spectrum of the M star companions of RE J0134-160 and RE J1126+183 are multiplied by 3.

Fig. 6.— Spectra of the magnetic white dwarfs PG 1658+441 and RE J1440+750. The position of the σ components of the Balmer lines are indicated.

Fig. 7.— Best fit of the magnetic white dwarf RE J1440+750. As indicated by the dashed line, $H\beta$ was not used for the parameter determination.

Fig. 8.— Differences in temperature (left), and gravity (right) between the M97, V97, and FKB samples and our own on a star by star basis. The same comparison is carried out for the LTE studies of M97 and V97 in the bottom panel. The smoothed average of the differences is plotted with solid lines, while the standard error of the mean difference is indicated by the dotted lines. The T_{eff} values used for the x-axis are NLTE from this paper (NGS), except for the bottom panel, which uses V97 (corrected to NLTE).

Fig. 9.— Effective temperature and gravity of our white dwarf sample compared with evolutionary tracks. Solid lines are Blöcker (1995); dashed lines are Driebe et al. (1998); dotted lines Wood (1995). The magnetic white dwarfs are marked by open symbols.

Fig. 10.— Mass distribution of white dwarfs (binned over $0.05 M_{\odot}$). The solid lined histogram indicates the result for our sample. The dotted line is the BSL sample with masses redetermined from the Blöcker/Driebe relations (normalized to the same sample size). The Gaussian curve represents the best fit to our mass distribution as described in the text.

Fig. 11.— Time evolution of stellar surface temperature for helium core white dwarfs (tracks from Driebe et al. 1998; solid lines) and C/O white dwarfs (tracks from Blöcker 1995; dashed lines). The tracks are labeled with the stellar masses.





















